

Perry H. Trester
D.M.D., F.R.C.D.(C), F.A.D.S.A., F.I.C.D., F.A.C.D.

Ali M.M. Sadeghi
D.M.D., M.D., F.R.C.D.(C)

H. Tony David
D.M.D., M.S.

Ben Kang
D.M.D., F.R.C.D.(C), Dip, A.B.O.M.S.

Joel B. Epstein
D.M.D., M.S.D., F.R.C.D.(C), Dip, A.B.O.M.



#209 - 179 Davie Street
Vancouver, BC V6Z 2Y1
Phone 604 688 7781
Fax 604 683 2203

vancouver@westcoastoralsurgery.ca
www.westcoastoralsurgery.ca

Bone Grafting Instructions

These instructions apply when grafting material has been placed into extraction sites to help preserve your jawbone, in preparation for possible implant replacement of the extracted tooth.

Your bone graft is made up of many particles. You may find some small granules in your mouth for the first several days. Do not be alarmed by these. It's normal to have some of them come out of the graft site and into your mouth.

There are some things you could do to minimize the amount of particles that become dislodged:

- *Do not disturb or touch the wound
- *Avoid rinsing or spitting for 2 days to allow blood clot and graft material stabilization
- *Do not apply pressure with your tongue or fingers to the grafted area, as the material is movable during the initial healing
- *Do not lift or pull on the lip to look at the sutures. This can actually cause damage to the wound site and tear the sutures
- *Do not smoke

Following the second day, gentle rinsing would be allowed, but not too vigorously, as you can again disturb some of the bone graft granules. If a partial denture or a flipper was placed in your mouth, you may have to see your restorative dentist to have it adjusted and learn how to remove and replace it appropriately.

It is our desire that your recovery be as smooth and pleasant as possible. Following these instructions will assist you, but if you have questions about your progress, please call the office.

After Sinus Lift

You have had a Sinus Lift Augmentation procedure in your upper jaw. This procedure regains lost bone height in the area of your first and second molar and occasionally second premolar. It is an important procedure as it allows implant placement in an area that could not be implanted otherwise, because of insufficient bone height. The bone that has been grafted is most commonly a combination of freeze-dried bone, artificial synthetic bone and your own bone.

Because of this you may have two post-surgical wounds: the donor site and the recipient site.

DO NOT UNDER ANY CIRCUMSTANCES BLOW YOUR NOSE FOR THE NEXT FOUR (4) WEEKS.

This may be longer if indicated. You may sniff all you like but NO BLOWING.

Do not blow your nose or sneeze holding your nose. Sneeze with your mouth open. Do not drink with straws and do not spit.

Scuba diving and flying in pressurized aircraft may also increase sinus pressure and should be avoided.

Decongestants such as Drixoral, Dimetapp, or Sudafed will help reduce pressure in the sinuses.

You will also be given a prescription for antibiotics. Please take these as directed.

Anything that causes pressure in your nasal cavity must be avoided. Avoid “bearing down”—as when lifting heavy objects, blowing up balloons, playing musical instruments that require a blowing action or any other activity that increases nasal or oral pressure.

Smoking must be stopped.

Antibiotics

Be sure to take the prescribed antibiotics as directed to help prevent infection.

Oral Hygiene

Do not rinse or spit on the day of your surgery. This tends to disturb the blood clot, open the wound and can prolong bleeding and slow healing.

You should not have a significant amount of blood in your mouth. Saliva can be swallowed, even if slightly blood tinged.

Keeping your mouth clean after surgery is essential to reduce the risk of infection. Start salt water rinses the day following your procedure. Use one-half teaspoon of salt dissolved in an 8 ounce glass of warm water and gently rinse with portions of the solution, taking five minutes to use the entire glassful. Repeat as often as you like, but at least four to five times daily and always after eating for the next five days.

Do not brush the teeth in the area of surgery for 48 hours. When brushing, be very gentle. When expectorating, also be gentle.

We may prescribe an antibiotic rinse (Chlorhexidine, Perio Plus, Peridex) for certain procedures. This rinse should be used in the morning and at bedtime after routine mouth care. Do not eat or drink or rinse your mouth after using the medicated rinse. Using this rinse more than two times a day will cause staining of your teeth and tongue. Make sure to brush your tongue also to minimize this staining.

Smoking

Do not smoke for at least two weeks after surgery, if at all. As discussed at your consultation, smoking dramatically increases the risk of bone graft and sinus augmentation failure. Your surgeon can prescribe a Nicoderm patch if you feel you need it.

Wearing your Denture or Nightguards:

Partial dentures, flippers, or full dentures should not be used immediately after surgery until your post-operative appointment unless specifically instructed otherwise. Please contact the office if there is any question. If you have questions about the fit of your flipper, partial or complete denture, do not wear it until your general dentist or our office can see you.

Post-Operative Problems or Complications

As with any procedure, unexpected post-operative healing can occur:

If you notice the unexpected flow of air or liquids between your mouth and nose, please let us know immediately

If you are aware of several small particles of graft material being discharged from your nose, let us know as well

If you experience sinus or nasal congestion on the side your surgery was performed, let us know

If there is an increase in swelling in your mouth, cheek or under your eye after 3 days, let us know

It is our desire that your recovery be as smooth and pleasant as possible. Following these instructions will assist you, but if you have questions about your progress, please call the office.